

SOUTH AFRICAN COMMITTEE OF MEDICAL DEANS

T: +27 (0) 82 930 6246 (Support and coordination) | +27 (0) 12 319 2192 (Chair)

E: sacomd@usaf.ac.za | P.O. Box 27392, Sunnyside, Pretoria, 0132 |

Care of: Universities South Africa, 1267 Pretorius Street, Block D, Hadefields Office Park, Hatfield, Pretoria, Gauteng, 0083

COMPULSORY COVID-19 VACCINATION RECOMMENDED FOR ALL HEALTH SCIENCES STUDENTS AND HEALTHCARE WORKERS

The South African Committee of Medical Deans (SACOMD) and the South African Committee of Dental Deans (SACODD) representing higher education institutions who educate the South African health workforce, recommend compulsory vaccination against the SARS-CoV-2 virus which causes Covid-19 for all health sciences students as well as the general health workforce. This will advance the efforts to curtail the spread of the virus amongst our communities and limit the impact the virus may have of significant disease. The further critical protection vaccination will afford, is for dental healthcare workers and students exposed to high level Aerosol Generating Procedures (AGPs).

We recognise the following scientific evidence which has emerged during the pandemic globally:

- the Covid-19 death rate in South Africa is 401 per 100,000 population as of mid-August 2021, placing it at the time among the top 10 countries with the highest Covid-19 death rate globally.
- Covid-19 vaccines have maintained high effectiveness against severe Covid-19, despite mutations affecting the spike protein.
- Covid-19 vaccination is estimated to lower the risk of transmission by 2-2.8 fold.
- Covid-19 vaccination halves the risk of long Covid-19 (symptoms for >28 days), which can be debilitating and affect the productivity of individuals.
- the higher level of SARS-CoV-2 exposure, including exposure to higher viral load, contributes to healthcare workers (HCW) being more susceptible to becoming infected, and possibly more likely to develop symptomatic and severe Covid-19. Studies from South African health facilities reported approximately 40% of HCWs were infected during the course of the first wave alone.

NELSON MANDELA
UNIVERSITY



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA



UNIVERSITY OF
KWAZULU-NATAL
INYUVESI
YAKWAZULU-NATALI



UNIVERSITY OF THE
FREE STATE
UNIVERSITEIT VAN DIE
VRYSTAAT
YUNIBESITHI YA
FREESTATA

UFS·UV
HEALTH SCIENCES
GESONDHEIDSWETENSKAPPE

- the higher risk of HCWs being infected by SARS-CoV-2, also increases the risk of their family members being infected by the virus. Household transmission studies of secondary cases resulting from HCWs being infected by SARS-CoV-2 report a 48% to 79% reduction in secondary cases in unvaccinated members in those households where HCWs had been vaccinated against Covid-19 (with two doses of the Pfizer or AstraZeneca vaccine) compared with where the index HCW cases were unvaccinated.
- HCWs may present additional risks to their patients as well as to the general households they come from, and their connected communities. Covid-19 vaccination of HCWs has the additional benefit of safeguarding healthcare services.

Historically, HCWs have been required as students to be vaccinated against Hepatitis B virus, albeit primarily for individual benefit in that case. The case for Covid-19 immunization transcends individual benefit, and has major benefits for the good of the broader health facility community, including fellow colleagues and patients whose lives are entrusted to the care of the HCW. As such, it is a moral imperative for HCWs to submit to vaccination against Covid-19, and for the best schedule of vaccine to be used to maximise protection not only against severe Covid-19, but also that which has the greatest likely impact in protecting against infection and mild-moderate Covid-19.

Our final recommendation is that all HCWs in South Africa who already received a single dose of the non-replicating vector J&J Covid-19 vaccine should receive a booster dose of Covid-19 vaccine. The single dose J&J vaccine in South African HCWs has been reported to reduce risk of hospitalization and death by 74% and 97%, respectively. However, data are unavailable of J&J effectiveness against infection and mild Covid-19, which is expected to be substantially lower than two doses of the Pfizer vaccine that induces substantially higher concentrations of neutralizing antibody than a single dose of J&J. A booster dose of J&J six months after vaccination has been shown to increase antibody levels nine-fold. Alternately, a heterologous prime-boost approach with a boost of a messenger RNA vaccine, for which evidence for the AstraZeneca non-replicating vector vaccine (which induces similar antibody responses even after a single dose compared with the J&J vaccine) followed by a single dose of the Pfizer vaccine exists, could be considered. The AZ-Pfizer heterologous schedule induces similar to greater neutralizing antibody and cell mediated immune responses than two doses of either of the vaccines, and particularly compared with two doses of the AZ vaccine.

SACOMD and SACODD make an urgent appeal through this statement for the following actions:

- 1) all adults in South Africa must be vaccinated against Covid-19 as soon as possible.
- 2) that Covid-19 vaccination be compulsory for all healthcare workers (HCWs) including those in training as health professions students in South Africa, also for the safety of patients and the general community.

- 3) serious consideration of booster doses for all HCWs in South Africa who already received a single dose of the non-replicating vector J&J Covid-19 vaccine.

PROF C. (TIAAN) DE JAGER

SACOMD CHAIR

E-mail: tiaan.dejager@up.ac.za

Tel: 012 319 2191/2192 or 083 703 1177